

Wang Y, Taylor PN, Baier G.

[Computational modelling of micro-seizures and focal seizure onset.](#)

*In: Twenty Second Annual Computational Neuroscience Meeting: CNS*2013.*

13-18 July 2013, Paris: BioMed Central.

Copyright:

© 2013 Wang et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Link to published item:

<http://dx.doi.org/10.1186/1471-2202-14-S1-P14>

Date deposited:

09/04/2015



This work is licensed under a [Creative Commons Attribution 2.0 Generic License](http://creativecommons.org/licenses/by/2.0)

POSTER PRESENTATION

Open Access

Computational modelling of micro-seizures and focal seizure onset

Yujiang Wang^{1*}, Peter N Taylor², Gerold Baier³

From Twenty Second Annual Computational Neuroscience Meeting: CNS*2013
Paris, France. 13-18 July 2013

Pathological micro-domains have been proposed to underpin the generation of local pathological activity, as seen in focal seizures in the epileptic cortex [1-3]. Specifically, so-called micro-seizures have been suggested to be markers for these micro-domains [2,3]. Astonishingly, micro-seizures have also been observed in non-epileptic

control patients [3]. This suggests that local activity, such as micro-seizures, only become pathological when in a specific arrangement.

We hypothesize that pathological dynamics could be due to an increased density of micro-domains. To test this, we introduce a computational model on the mesoscopic scale




Figure 1 First column indicated the location of the pathological clusters in black and the position of the micro-incision using the labeled line. The second column shows the temporal average activity in 10000 simulation steps. The third column indicates the spatial average of the same simulation using the black line and the purple lines show the spatial average of 10×10 subsquares, which represent macro-columns in the model. (A) Resulting simulation after a 5 minicolumn long micro-incision. (B) Resulting simulation after a 10 minicolumn long micro-incision. (C) Resulting simulation after a 45 minicolumn long micro-incision.

* Correspondence: yujiang.wang@manchester.ac.uk

¹Manchester Interdisciplinary Biocentre, 131 Princess Street, Manchester M1 7DN, UK

Full list of author information is available at the end of the article

of a $5 \times 5 \text{ mm}^2$ cortical sheet [4]. The units are modelled as excitable minicolumns. This model also incorporates realistic connectivity schemes observed at this spatial scale [5].

The model shows occasional, non-pathological micro-seizure occurrences, as well as recruitment of normal tissue into full-blown seizure activity in the presence of dense clusters of hyperactive micro-seizure domains. A specific prediction of this model is that the transition to full-blown seizures can be prevented by using micro-incisions to separate the clusters of abnormally active micro-domains (Figure. 1) [6].

Author details

¹Manchester Interdisciplinary Biocentre, 131 Princess Street, Manchester M1 7DN, UK. ²School of Electrical & Electronic Engineering, Nanyang Technological University, Singapore. ³Division of Bioscience, Faculty of Life Sciences, University College London, London WC1E 6BT, UK.

Published: 8 July 2013

References

1. Bragin A, Wilson CL, Engel J Jr: **Chronic epileptogenesis requires development of a network of pathologically interconnected neuron clusters: a hypothesis.** *Epilepsia* 2000, **41**:144-152.
2. Schevon CA, Ng SK, Cappell J, Goodman RR, McKhann G, Waziri A, Branner A, Sosunov A, Schroeder CE, Emerson RG: **Microphysiology of Epileptiform Activity in Human Neocortex.** *J Clin Neurophysiol* 2008, **25**:321-330.
3. Stead M, Bower M, Brinkmann BH, Lee K, Marsh WR, Meyer FB, Litt B, Van Gompel J, Worrell GA: **Microseizures and the spatiotemporal scales of human partial epilepsy.** *Brain* 2010, **133**:2789-2797.
4. Wang Y, Goodfellow M, Taylor PN, Gary DJ, Baier G: **Computational Modelling of Microseizures and Focal Seizure Onset.** In *Proceedings of IWSP5*. World Scientific; Lehnertz, R, Tetzlaff, C, Elger 2013.
5. Voges N, Schüz A, Aertsen A, Rotter S: **A modeler's view on the spatial structure of intrinsic horizontal connectivity in the neocortex.** *Prog Neurobiol* 2010, **92**:277-292.
6. Wang Y, Goodfellow M, Taylor P, Garry D, Baier G: **Computational modelling of microseizures and focal seizure onset.** *International Seizure prediction workshop proceedings*, Dresden 2011 (to appear).

doi:10.1186/1471-2202-14-S1-P14

Cite this article as: Wang et al.: Computational modelling of micro-seizures and focal seizure onset. *BMC Neuroscience* 2013 **14**(Suppl 1):P14.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

